




Lagrange Multipliers



Lagrange Multipliers

Many optimization problems have restrictions, or **constraints**, on the values that can be used to produce the optimal solution. Such constraints tend to complicate optimization problems because the optimal solution can occur at a boundary point of the domain. In this section, you will study an ingenious technique for solving such problems. It is called the **Method of Lagrange Multipliers**.

To see how this technique works, suppose you want to find the rectangle of maximum area that can be inscribed in the ellipse given by

$$\frac{x^2}{3^2} + \frac{y^2}{4^2} = 1.$$

Let (x, y) be the vertex of the rectangle in the first quadrant, as shown in Figure 13.77. Because the rectangle has sides of lengths $2x$ and $2y$, its area is given by

$$f(x, y) = 4xy. \quad \text{Objective function}$$

You want to find x and y such that $f(x, y)$ is a maximum. Your choice of (x, y) is restricted to first-quadrant points that lie on the ellipse

$$\frac{x^2}{3^2} + \frac{y^2}{4^2} = 1. \quad \text{Constraint}$$

Now, consider the constraint equation to be a fixed level curve of

$$g(x, y) = \frac{x^2}{3^2} + \frac{y^2}{4^2}.$$

The level curves of f represent a family of hyperbolas

$$f(x, y) = 4xy = k.$$

In this family, the level curves that meet the given constraint correspond to the hyperbolas that intersect the ellipse. Moreover, to maximize $f(x, y)$, you want to find the hyperbola that just barely satisfies the constraint. The level curve that does this is the one that is *tangent* to the ellipse, as shown in Figure 13.78.





Fig.13.77




Ellipse:
 $\frac{x^2}{3^2} + \frac{y^2}{4^2} = 1$



Objective function: $f(x, y) = 4xy$

Level curves of f :
 $4xy = k$



Constraint: $g(x, y) = \frac{x^2}{3^2} + \frac{y^2}{4^2} = 1$



Recall




Fig. 13.48


The Gradient of a Function of Two Variables

Definition of Gradient of a Function of Two Variables

Let $z = f(x, y)$ be a function of x and y such that f_x and f_y exist. Then the **gradient of f** , denoted by $\nabla f(x, y)$, is the vector

$$\nabla f(x, y) = f_x(x, y)\mathbf{i} + f_y(x, y)\mathbf{j}.$$

∇f is read as “del f .” Another notation for the gradient is **grad** $f(x, y)$. In Figure 13.48, note that for each (x, y) , the gradient $\nabla f(x, y)$ is a vector in the plane (not a vector in space).



To find the appropriate hyperbola, use the fact that two curves are tangent at a point if and only if their gradient vectors are parallel. This means that $\nabla f(x, y)$ must be a scalar multiple of $\nabla g(x, y)$ at the point of tangency. In the context of constrained optimization problems, this scalar is denoted by λ (the lowercase Greek letter lambda).


$$\nabla f(x, y) = \lambda \nabla g(x, y)$$

The scalar λ is called a **Lagrange multiplier**. Theorem 13.19 gives the necessary conditions for the existence of such multipliers.

THEOREM 13.19 Lagrange's Theorem

Let f and g have continuous first partial derivatives such that f has an extremum at a point (x_0, y_0) on the smooth constraint curve $g(x, y) = c$. If $\nabla g(x_0, y_0) \neq \mathbf{0}$, then there is a real number λ such that

$$\nabla f(x_0, y_0) = \lambda \nabla g(x_0, y_0).$$



The Method of Lagrange Multipliers uses Theorem 13.19 to find the extreme values of a function f subject to a constraint.

Method of Lagrange Multipliers

Let f and g satisfy the hypothesis of Lagrange's Theorem, and let f have a minimum or maximum subject to the constraint $g(x, y) = c$. To find the minimum or maximum of f , use the following steps.



1. Simultaneously solve the equations $\nabla f(x, y) = \lambda \nabla g(x, y)$ and $g(x, y) = c$ by solving the following system of equations.


$$f_x(x, y) = \lambda g_x(x, y)$$

$$f_y(x, y) = \lambda g_y(x, y)$$


$$g(x, y) = c$$

2. Evaluate f at each solution point obtained in the first step. The largest value yields the maximum of f subject to the constraint $g(x, y) = c$, and the smallest value yields the minimum of f subject to the constraint $g(x, y) = c$.






Use Lagrange
Multipliers to solve
problems



Problem: Suppose I have 50 feet of fencing for a rectangular shaped garden and want to enclose the maximum area.

$$2x + 2y = 50 \text{ Constraint}$$

$$A = xy \text{ Objective}$$



x

We want to get the Area in terms of just x or y so we use the constraint to eliminate one or the other.

$$2y = 50 - 2x$$

$$y = 25 - x$$

$$A = x(25 - x) = 25x - x^2$$

$$A' = 25 - 2x$$

$$0 = 25 - 2x$$

$$x = 12.5 \text{ and } y = 12.5$$



Now we will use the method of **Lagrange Multipliers** to Solve the same problem.

$$2x + 2y = 50 \text{ Constraint}$$

$$A = xy \text{ Objective}$$

$f(x,y)=xy$ is the objective equation and $g(x,y)=2x +2y$ is the constraint equation.

Find $f_x(x, y) = y; f_y(x, y) = x$

$$g_x(x, y) = 2; g_y(x, y) = 2$$

$$\begin{cases} \text{Set } f_x = \lambda g_x \rightarrow y = 2\lambda \\ \text{Set } f_y = \lambda g_y \rightarrow x = 2\lambda \\ 2x + 2y = 50 \end{cases}$$

Solve simultaneously


$$4\lambda + 4\lambda = 50$$

$$8\lambda = 50$$

$$\lambda = 6.25$$

$$x = 12.5$$

$$y = 12.5$$



Now let's look at a problem that might be difficult to solve the old way.

Problem 2: Find the maximum and minimum values for

$$f(x, y) = x^2 + y^2 \text{ objective equation}$$

$$g(x, y) = x^4 + y^4 \text{ with the constraint } x^4 + y^4 = 1$$

$$\text{Find } f_x = 2x ; f_y = 2y$$

$$g_x = 4x^3 ; g_y = 4y^3$$

$$\begin{cases} 2x = \lambda 4x^3 \\ 2y = \lambda 4y^3 \\ x^4 + y^4 = 1 \end{cases}$$

$$2x - \lambda 4x^3 = 2x(1 - \lambda 2x^2) \Rightarrow x = 0 \text{ or } \lambda = \frac{1}{2x^2}$$

$$2y - \lambda 4y^3 = 2y(1 - \lambda 2y^2) \Rightarrow y = 0 \text{ or } \lambda = \frac{1}{2y^2}$$

$$\text{Solve: } \frac{1}{2x^2} = \frac{1}{2y^2} \Rightarrow x^2 = y^2$$


$$\text{If } x = 0 \text{ then } y^4 = 1 \Rightarrow y = \pm 1$$


$$x^4 + y^4 = 1 \Rightarrow x^4 + x^4 = 1 \Rightarrow 2x^4 = 1$$

$$\text{If } y = 0 \text{ then } x^4 = 1 \Rightarrow x = \pm 1$$

$$x = \frac{1}{\sqrt[4]{2}} \text{ and } y = \frac{1}{\sqrt[4]{2}}$$

We could have the following points $\left(\pm \frac{1}{\sqrt[4]{2}}, \pm \frac{1}{\sqrt[4]{2}} \right)$ which yield a max value of $\sqrt{2}$ and $(0, \pm 1), (\pm 1, 0)$ which yields a min value of 1





The constraint is the red graph.

The blue graph and green graph are level curves of the paraboloid.

The blue graph is at 1 and the green graph is at $\sqrt{2}$

which is the exact value for the max.


As the paraboloid extends in the z direction it gets wider than the constraint.



Suppose we want to find the max and min values of $f(x,y,z)$ subject to **two constraints** of the form $g(x,y,z)=p$ and $h(x,y,z)=q$. Geometrically, this means that we are looking for the extreme values of f when (x,y,z) is restricted to lie on the curve of intersection of the level surfaces g and h . It can be shown that if an extreme value occurs at (x_o, y_o, z_o) , then the gradient vector $\nabla f(x_o, y_o, z_o)$ is in the plane determined by $\nabla g(x_o, y_o, z_o)$ and $\nabla h(x_o, y_o, z_o)$.

We assume these gradient vectors are not 0 or parallel and thus there are numbers λ and μ (Lagrange multipliers) such that

$$\nabla f(x_o, y_o, z_o) = \lambda \nabla g(x_o, y_o, z_o) + \mu \nabla h(x_o, y_o, z_o)$$



Problem: Maximize $f(x,y,z) = xyz$ subject to the two constraints $x^2 + z^2 = 5$ and $x - 2y = 0$

$$f(x, y, z) = xyz \quad f_x = yz \quad f_y = xz \quad f_z = xy$$


$$g(x, y, z) = x^2 + z^2 \quad g_x = 2x \quad g_y = 0 \quad g_z = 2z$$

$$h(x, y, z) = x - 2y \quad h_x = 1 \quad h_y = -2 \quad h_z = 0$$

$$x^2 + z^2 = 5 \text{ and } x - 2y = 0$$

Solve

$$\begin{cases} yz = \lambda 2x + \mu \\ xz = \lambda 0 - \mu 2 \\ xy = \lambda 2z + \mu 0 \\ x^2 + z^2 = 5 \text{ and } x - 2y = 0 \end{cases}$$



$$xz = -2\mu \Rightarrow \mu = \frac{-xz}{2}$$

$$xy = \lambda 2z \Rightarrow \lambda = \frac{xy}{2z}$$

$$yz = \frac{xy}{2z} 2x + \frac{-xz}{2}$$

$$2yz^2 = 2x^2y - xz^2$$

Sub $y = \frac{x}{2}$ from $x - 2y = 0$

Sub $z^2 = 5 - x^2$ from $x^2 + z^2 = 5$

$$2 \frac{x}{2} (5 - x^2) = 2x^2 \frac{x}{2} - x(5 - x^2)$$

$$x(5 - x^2) = x^2x - x(5 - x^2)$$

$$2x(5 - x^2) = x^3$$

$$10x - 2x^3 - x^3 = 0$$

$$10x - 3x^3 = 0$$

$$x(10 - 3x^2) = 0$$

$$x = 0, \text{ or } x^2 = \frac{10}{3} \Rightarrow x = \pm \sqrt{\frac{10}{3}}$$

Continued

$$y = \frac{x}{2}$$

$$z^2 = 5 - x^2$$

$$x = 0, \text{ or } x^2 = \frac{10}{3} \Rightarrow x = \pm \sqrt{\frac{10}{3}}$$

for $x = 0, y = 0, z = \pm \sqrt{5}$

for $x = \sqrt{\frac{10}{3}}, y = \frac{1}{2} \sqrt{\frac{10}{3}},$

$$z^2 = 5 - \frac{10}{3} = \frac{5}{3}, z = \pm \sqrt{\frac{5}{3}}$$

points

$$(0, 0, \sqrt{5}),$$

$$\left(\sqrt{\frac{10}{3}}, \frac{1}{2} \sqrt{\frac{10}{3}}, \sqrt{\frac{5}{3}} \right)$$

$$\left(-\sqrt{\frac{10}{3}}, -\frac{1}{2} \sqrt{\frac{10}{3}}, \sqrt{\frac{5}{3}} \right)$$

$$f(0, 0, \sqrt{5}) = 0$$

$$f\left(\sqrt{\frac{10}{3}}, \frac{1}{2} \sqrt{\frac{10}{3}}, \sqrt{\frac{5}{3}} \right) = \frac{5\sqrt{15}}{9}$$

$$f\left(-\sqrt{\frac{10}{3}}, -\frac{1}{2} \sqrt{\frac{10}{3}}, \sqrt{\frac{5}{3}} \right) = \frac{5\sqrt{15}}{9}$$

