

# Vector spaces

Vector spaces, subspaces & basis

# A vector space $(V, F, +, \cdot)$

- $F$  a field
- $V$  a set (of objects called vectors)
- **Addition** of vectors (commutative, associative)
- Scalar **multiplication**  $\exists 0, \forall \alpha \in V, \alpha + 0 = 0$ .

$$\forall \alpha \exists! -\alpha, \alpha + (-\alpha) = 0.$$

$$(c, \alpha) \mapsto c\alpha, c \in F, \alpha \in V$$

$$1\alpha = \alpha, (c_1 c_2)\alpha = c_1(c_2\alpha), c(\alpha + \beta) = c\alpha + c\beta, (c_1 + c_2)\alpha = c_1\alpha + c_2\alpha$$

# Examples

$$F^n = \{(x_1, \dots, x_n) \mid x_i \in F\}$$

$$\begin{aligned}(x_1, \dots, x_n) + (y_1, \dots, y_n) &= (x_1 + y_1, \dots, x_n + y_n) \\ c(x_1, \dots, x_n) &= (cx_1, \dots, cx_n)\end{aligned}$$

- Other laws are easy to show

$$\mathbf{C}^n, (Q + \sqrt{2}Q)^n, \mathbf{Z}_p^n$$

$$F^{m \times n} = \{\{A_{ij}\} \mid A_{ij} \in F, i=1, \dots, m, j=1, \dots, n\} =$$

$$F^{mn} = \{(A_{11}, A_{12}, \dots, A_{m-1,1}, A_{m-1,m}) \mid A_{ij} \in F\}$$

- This is just written differently



- The space of functions:  $A$  a set,  $F$  a field

$$\{f : A \rightarrow F\}, (f + g)(s) = f(s) + g(s), (cf)(s) = c(f(s))$$

- If  $A$  is finite, this is just  $F^{|A|}$ . Otherwise this is infinite dimensional.

- The space of polynomial functions

$$\{f : F \rightarrow F \mid f(x) = c_0 + c_1x + c_2x^2 + \dots + c_nx^n, c_i \in F\}$$

- The following are different.

$$\begin{array}{l} V = \mathbf{C} = \{x + iy \mid x, y \in \mathbf{R}\} \quad , \quad F = \mathbf{R} \\ \quad \quad \quad V = \mathbf{C} \quad \quad \quad , \quad F = \mathbf{C} \\ \quad \quad \quad V = \mathbf{C} \quad \quad \quad , \quad F = \mathbf{Q} \end{array}$$

# Subspaces

- $V$  a vector space of a field  $F$ . A **subspace**  $W$  of  $V$  is a subset  $W$  s.t. restricted operations of vector addition, scalar multiplication make  $W$  into a vector space.
  - $+:W \times W \rightarrow W$ ,  $\cdot:F \times W \rightarrow W$ .
  - $W$  nonempty subset of  $V$  is a vector subspace iff for each pair of vectors  $a, b$  in  $W$ , and  $c$  in  $F$ ,  $ca+b$  is in  $W$ .  
(iff for all  $a, b$  in  $W$ ,  $c, d$  in  $F$ ,  $ca+db$  is in  $W$ .)
- Example:

$$\mathbf{R}^{n-1} \subset \mathbf{R}^n, \{(x_1, \dots, x_{n-1}, 0) \mid x_i \in \mathbf{R}\}$$



- $S_{m \times m} = \{A \in F^{m \times m} \mid A^t = A\} \subset F^{m \times m}$   
is a vector subspace with field F.
- **Solution spaces:** Given an  $m \times n$  matrix A

$$W = \{X \in F^n \mid AX = 0\} \subset F^n$$

$$\forall X, Y \in W, c \in F, A(cX + Y) = cAX + AY = 0. \mapsto cX + Y \in W.$$

- The intersection of a collection of vector subspaces is a vector subspace
- $\bigcup_{i=1}^n W_i$  is not.

$$W = \{(x, y, z) \mid x = 0 \text{ or } y = 0\}$$

# Span(S)

$$\text{Span}(S) = \left\{ \sum_i c_i \alpha_i \mid \alpha_i \in S, c_i \in F \right\} \subset V$$

- Theorem 3.  $W = \text{Span}(S)$  is a vector subspace and is the set of all linear combinations of vectors in  $S$ .
- Proof:

$$c = \sum_{i=1}^m c_i \alpha_i + \sum_{j=1}^n d_j \beta_j$$

$$= X_{1 \ 1} + \cdots + X_{m \ m}$$

$$= y_{1 \ 1} + \cdots + y_{n \ n}$$

$$c + \quad = X_{1 \ 1} + \cdots + X_{m \ m} + y_{1 \ 1} + \cdots + y_{n \ n}$$



- **Sum of subsets**  $S_1, S_2, \dots, S_k$  of  $V$

$$S_1 + S_2 + \dots + S_k = \{\alpha_1 + \alpha_2 + \dots + \alpha_k \mid \alpha_i \in S_i\}$$

- If  $S_i$  are all subspaces of  $V$ , then the above is a subspace.
- Example:  $y=x+z$  subspace:

$$\text{Span}((1, 1, 0), (0, 1, 1)) = \{c(1, 1, 0) + d(0, 1, 1) \mid c, d \in \mathbf{R}\} = \{(c, c+d, d) \mid c, d \in \mathbf{R}\}$$

- Column space of  $A$ : the space of column vectors of  $A$ .



# Linear independence

- A subset  $S$  of  $V$  is **linearly dependent** if

$\exists \alpha_1, \dots, \alpha_n \in S, c_1, \dots, c_n \in F$  not all 0 s.t.  $c_1\alpha_1 + \dots + c_n\alpha_n = 0$ .

- A set which is not linearly dependent is called **linearly independent**:

The negation of the above statement

$\forall \alpha_1, \dots, \alpha_n \in S$ , there are no  $c_1, \dots, c_n \in F$  not all 0 such that  $c_1\alpha_1 + \dots + c_n\alpha_n = 0$ .)

$\forall \alpha_1, \dots, \alpha_n \in S$ , if  $c_1\alpha_1 + \dots + c_n\alpha_n = 0$ , then  $c_i = 0, i = 1, \dots, n$

$$(1, 1), (0, 1), c_1(1, 1) + c_2(0, 1) = (c_1, c_1 + c_2) = (0, 0) \mapsto c_1 = 0, c_2 = 0$$

$$c_1(1, 1, 1) + c_2(2, 2, 1) + c_3(3, 3, 2) = 0 \text{ for } c_1 = 1, c_2 = 1, c_3 = -1.$$

# Basis

- A **basis** of  $V$  is a linearly independent set of vectors in  $V$  which spans  $V$ .
- Example:  $F^n$  the standard basis  
 $\epsilon_1 = (1, 0, \dots, 0), \epsilon_2 = (0, 1, \dots, 0), \dots, \epsilon_n = (0, 0, \dots, 1)$
- $V$  is **finite dimensional** if there is a finite basis.  
**Dimension** of  $V$  is the number of elements of a basis. (Independent of the choice of basis.)
- A proper subspace  $W$  of  $V$  has  $\dim W < \dim V$ . (to be proved)



- Example:  $P$  invertible  $n \times n$  matrix.  $P_1, \dots, P_n$  columns form a basis of  $F^{n \times 1}$ .
  - Independence:  $x_1 P_1 + \dots + x_n P_n = 0$ ,  $PX = 0$ . Thus  $X = 0$ .
  - Span  $F^{n \times 1}$ :  $Y$  in  $F^{n \times 1}$ . Let  $X = P^{-1}Y$ . Then  $Y = PX$ .  $Y = x_1 P_1 + \dots + x_n P_n$ .
- Solution space of  $AX = 0$ . Change to  $RX = 0$ .

$$\begin{array}{rcl}
 x_{k_1} & + & \sum_{j=1}^{n-r} C_{1j} u_j = 0 \\
 & & x_{k_2} + \sum_{j=1}^{n-r} C_{2j} u_j = 0 \\
 & & \ddots + \vdots = \vdots \\
 & & x_{k_r} + \sum_{j=1}^{n-r} C_{rj} u_j = 0
 \end{array}$$

- Basis  $E_j$   $u_j = 1$ , other  $u_k = 0$  and solve above

$$x_{k_i} = -C_{ij}, \mapsto (-C_{1j}, -C_{2j}, \dots, -C_{rj}, 0, \dots, 1, \dots, 0)$$



- Thus the dimension is  $n-r$ :

- Infinite dimensional example:

- $V := \{f \mid f(x) = c_0 + c_1x + c_2x^2 + \dots + c_nx^n\}$ .

- Given any finite collection  $g_1, \dots, g_n$  there is a maximum degree  $k$ . Then any polynomial of degree larger than  $k$  can not be written as a linear combination.

- **Theorem 4:**  $V$  is spanned by  $\beta_1, \beta_2, \dots, \beta_m$   
Then any independent set of vectors in  $V$  is finite and number is  $\leq m$ .

- Proof: To prove, we show every set  $S$  with more than  $m$  vectors is linearly dependent. Let  $\alpha_1, \alpha_2, \dots, \alpha_n$  be elements of  $S$  with  $n > m$ .

$$\alpha_j = \sum_{i=1}^m A_{ij} \beta_i$$

$$\sum_{i=1}^n x_j \alpha_j = \sum_{j=1}^n x_j \sum_{i=1}^m A_{ij} \beta_i = \sum_{i=1}^m \left( \sum_{j=1}^n A_{ij} x_j \right) \beta_i$$

- $A$  is  $m \times n$  matrix. Theorem 6, Ch 1, we can solve for  $x_1, x_2, \dots, x_n$  not all zero for

$$\sum_{j=1}^n A_{ij} x_j = 0, i = 1, \dots, m$$

- Thus

$$x_1 \alpha_1 + \dots + x_n \alpha_n = 0$$



- **Corollary.**  $V$  is a finite d.v.s. Any two bases have the same number of elements.

- Proof:  $B, B'$  basis. Then  $|B'| \leq |B|$  and  $|B| \leq |B'|$ .

- This defines **dimension**.

- $\dim F^n = n$ .  $\dim F^{m \times n} = mn$ .

- **Lemma.**  $S$  a linearly independent subset of  $V$ . Suppose that  $b$  is a vector not in the span of  $S$ . Then  $S \cup \{b\}$  is independent.

- Proof: 
$$c_1\alpha_1 + \cdots + c_m\alpha_m + kb = 0.$$
 Then  $k=0$ . Otherwise  $b$  is in the span.

Thus,

and  $c_i$  are all zero. 
$$c_1\alpha_1 + \cdots + c_m\alpha_m = 0.$$



- **Theorem 5.** If  $W$  is a subspace of  $V$ , every linearly independent subset of  $W$  is finite and is a part of a basis of  $W$ .
- $W$  a subspace of  $V$ .  $\dim W \leq \dim V$ .
- A set of linearly independent vectors can be extended to a basis.
- A  $n \times n$ -matrix. Rows (respectively columns) of  $A$  are independent iff  $A$  is invertible.
  - ( $\rightarrow$ ) Rows of  $A$  are independent.  $\dim \text{Rows } A = n$ .  $\dim \text{Rows r.r.e } R$  of  $A = n$ .  $R$  is  $I \rightarrow A$  is inv.
  - ( $\leftarrow$ )  $A = B \cdot R$ . for r.r.e form  $R$ .  $B$  is inv.  $AB^{-1}$  is inv.  $R$  is inv.  $R = I$ . Rows of  $R$  are independent.  $\dim \text{Span } R = n$ .  $\dim \text{Span } A = n$ . Rows of  $A$  are independent.

- **Theorem 6.**

$$\dim (W_1+W_2) = \dim W_1+\dim W_2-\dim W_1\cap W_2.$$

- **Proof:**

- $W_1\cap W_2$  has basis  $a_1,\dots,a_k$ .  $W_1$  has basis  $a_1,\dots,a_k,b_1,\dots,b_m$ .  $W_2$  has basis  $a_1,\dots,a_k,c_1,\dots,c_n$ .
- $W_1+W_2$  is spanned by  $a_1,\dots,a_k,b_1,\dots,b_m,c_1,\dots,c_n$ .
- There are also independent.

- Suppose

$$\sum_{i=1}^l x_i a_i + \sum_{j=1}^m y_j b_j + \sum_{k=1}^n z_k c_k = 0$$

- Then

$$\sum_{k=1}^n z_k c_k = -\sum_{i=1}^l x_i a_i - \sum_{j=1}^m y_j b_j$$

$$\sum_{k=1}^n z_k c_k \in W_1 \text{ and } \in W_2$$

$$\sum_{k=1}^n z_k c_k = \sum_{i=1}^l d_i a_i$$

- By independence  $z_k=0$ .  $x_i=0,y_j=0$  also.



# Coordinates

- Given a vector in a vector space, how does one name it? Think of charting earth.
- If we are given  $F^n$ , this is easy? What about others?
- We use ordered basis:  
One can write any vector uniquely

$$\mathcal{B} = \{\alpha_1, \dots, \alpha_n\}$$

$$\alpha = x_1\alpha_1 + \dots + x_n\alpha_n$$



- Thus, we name

$$\alpha \mapsto (x_1, \dots, x_n) \in F^n \quad X = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = [\alpha]_{\mathcal{B}}$$

**Coordinate (n×1)-matrix (n-tuple) of a vector.**

For standard basis in  $F^n$ , coordinate and vector are the same.

- This sets up a **one-to-one correspondence** between  $V$  and  $F^n$ .
  - Given a vector, there is unique n-tuple of coordinates.
  - Given an n-tuple of coordinates, there is a unique vector with that coordinates.
  - These are verified by the properties of the notion of bases. (See page 50)

# Coordinate change?

- If we choose different basis, what happens to the coordinates?
- Given two bases

- Write

$$\mathcal{B} = \{\alpha_1, \dots, \alpha_n\}, \mathcal{B}' = \{\alpha'_1, \dots, \alpha'_n\}$$

$$\alpha'_j = \sum_{i=1}^n P_{ij} \alpha_i$$

$$\begin{aligned} \alpha &= \sum_{j=1}^n x_j \alpha_j = x_1 \alpha_1 + \dots + x_n \alpha_n \\ &= \sum_{j=1}^n x'_j \alpha'_j = \sum_{j=1}^n x'_j \sum_{i=1}^n P_{ij} \alpha_i \\ &= \sum_{j=1}^n \sum_{i=1}^n (P_{ij} x'_j) \alpha_i = \sum_{i=1}^n \left( \sum_{j=1}^n P_{ij} x'_j \right) \alpha_i. \\ x_i &= \sum_{j=1}^n P_{ij} x'_j \end{aligned}$$



- $X=0$  iff  $X'=0$  Theorem 7, Ch1,  $P$  is invertible

- Thus,  $X = PX'$ ,  $X'=P^{-1}X$ .

$$[\alpha]_{\mathcal{B}} = P[\alpha]_{\mathcal{B}'}, [\alpha]_{\mathcal{B}'} = P^{-1}[\alpha]_{\mathcal{B}},$$

- Example  $\{(1,0), (0,1)\}$ ,  $\{(1,i), (i,1)\}$

- $(1,i) = (1,0) + i(0,1)$

- $(i,1) = i(1,0) + (0,1)$

$$P = \begin{pmatrix} 1 & i \\ i & 1 \end{pmatrix}, P^{-1} = \begin{pmatrix} 1/2 & -i/2 \\ -i/2 & 1/2 \end{pmatrix},$$

- $(a,b) = a(1,0) + b(0,1)$ :  $(a,b)_{\mathcal{B}} = (a,b)$

- $(a,b)_{\mathcal{B}'} = P^{-1}(a,b) = ((a-ib)/2, (-ia+b)/2)$ .

- We check that  $(a-ib)/2x(1,i) + (-ia+b)/2x(i,1) = (a,b)$ .