

Cauchy-Riemann Equation

Functions of a complex variable

Let S be a set of complex numbers.

A function defined on S is a rule that assigns to each z in S a complex number w .

value of f at z , or $f(z)$

or

$$w = f(z)$$

S is the domain of definition of f

$w = \frac{1}{z}$ sometimes refer to the function f itself, for simplicity.

$$w = z^2 + 1$$

Both a domain of definition and a rule are needed in order for a function to be well defined.

Suppose $w = u + iv$ is the value of a function at f $z = x + iy$

$$u + iv = f(x + iy)$$

$$\text{or } f(z) = u(x, y) + iv(x, y)$$


real-valued functions of real variables x, y

$$\text{or } f(z) = u(r, \theta) + iv(r, \theta)$$

Ex.

$$f(z) = z^2$$

$$f(x + iy) = x^2 - y^2 + i2xy$$

$$u(x, y) = x^2 - y^2, \quad v(x, y) = 2xy$$

$$f(re^{i\theta}) = r^2 \cos 2\theta + ir^2 \sin 2\theta$$

$$u(r, \theta) = r^2 \cos 2\theta \quad v(r, \theta) = r^2 \sin 2\theta$$

when $v=0$

$f(z)$ is a real-valued function of a complex variable.

$f(z) = P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n$ is a polynomial of degree n .

$\frac{P(z)}{Q(z)}$: rational function, defined when $Q(z) \neq 0$

For multiple-valued functions : usually assign one to get single-valued function

Ex. $z = re^{i\theta}, \quad z \neq 0$

$$z^{\frac{1}{2}} = \pm \sqrt{r}e^{i\theta/2}, \quad -\pi < \theta \leq \pi \quad \text{nth root}$$

If we choose $f(z) = \sqrt{r}e^{i\theta/2} \quad (r > 0, -\pi < \theta < \pi)$

$$-\frac{\pi}{2} < \frac{\theta}{2} < \frac{\pi}{2}$$

and $f(0) = 0$,


then f is well defined on the entire complex plane except the ray $\theta = \pi$.

Mappings

$w=f(z)$ is not easy to graph as real functions are.

One can display some information about the function by indicating pairs of corresponding points $z=(x,y)$ and $w=(u,v)$. (draw z and w planes separately).

When a function f is thought of in this way, it is often referred to as a mapping, or transformation.



Mapping can be translation, rotation, reflection. In such cases it is convenient to consider z and w planes to be the same.

$$w = z + 1 \quad \text{translation} \quad +1$$

$$w = iz \quad \text{rotation} \quad \theta$$


$$w = \bar{z} \quad \text{reflection in real axis.} \quad \frac{\theta}{2}$$

Ex. image of curves


$$w = z^2$$

$$u = x^2 - y^2$$


$$v = 2xy$$



a hyperbola $x^2 - y^2 = c_1$ is mapped in a one to one manner onto the line $u = c_1$





right hand branch $x > 0$,
 left hand branch $x < 0$



image

$$\begin{aligned}
 u = C_1, \quad & V = 2y\sqrt{y^2 + c_1} \quad (-\infty < y < \infty) \\
 u = C_1, \quad & V = -2y\sqrt{y^2 + c_1} \quad (-\infty < y < \infty)
 \end{aligned}$$

Ex 2.



When $0 < x_1 < 1$, point (x_1, y) moves up a vertical half line, L_1 , as y increases from $y = 0$.

$$u = x^2 - y^2, \quad v = 2x_1 y$$

$$y = \frac{v}{2x_1}$$

$$u = x_1^2 - \left(\frac{v}{2x_1}\right)^2, \quad v^2 = -4x_1^2(u - x_1^2) \quad \leftarrow \text{ a parabola with vertex at } (x_1^2, 0)$$

half line CD is mapped of half line C'D'

$$(0, y)$$


$$(-y^2, 0)$$

Ex 3.


$$w = z^2 = r^2 e^{i2\theta}$$

$$\text{let } w = \rho e^{i\phi}$$

$$\rho = r^2, \quad \phi = 2\theta + 2n\pi \quad (n = 0, \pm 1, \pm 2, \dots)$$



$$r \geq 0, \quad 0 \leq \theta \leq \pi/2$$



one to one \longrightarrow $\rho \geq 0, \quad 0 \leq \phi \leq \pi$

Limits

Let a function f be defined at all points z in some deleted neighborhood of z_0



$$\lim_{z \rightarrow z_0} f(z) = w_0 \quad (1)$$

means: the limit of $f(z)$ as z approaches z_0 is w_0

$w = f(z)$ can be made arbitrarily close to w_0 if we choose the point z close enough to z_0 but distinct from it.

(1) means that, for each positive number ϵ , there is a positive number δ such that

$$|f(z) - w_0| < \epsilon \quad \text{whenever} \quad 0 < |z - z_0| < \delta \quad (2)$$



Note:

(2) requires that f be defined at all points in some deleted neighborhood of z_0

such a deleted neighborhood always exists when z_0 is an interior point of a region on which f is defined. We can extend the definition of limit to the case in which z_0 is a boundary point of the region by agreeing that left of (2) be satisfied by only those points z that lie in both the region and the domain

$0 < |z - z_0| < \delta$
Example 1. show if

$$f(z) = \frac{iz}{2} \quad \text{in } |z| < 1, \quad \text{then}$$


$$\lim_{z \rightarrow 1} f(z) = \frac{i}{2}$$

when z in $|z| < 1$

$$\left| f(z) - \frac{i}{2} \right| = \left| \frac{iz}{2} - \frac{i}{2} \right| = \frac{|z-1|}{2}$$

For any such z and any positive number ε

$$\left| f(z) - \frac{i}{2} \right| < \varepsilon \quad \text{whenever} \quad 0 < |z-1| < 2\varepsilon$$



When a limit of a function $f(z)$ exists at a point z_0 , it is unique.

If not, suppose $\lim_{z \rightarrow z_0} f(z) = w_0$ and $\lim_{z \rightarrow z_0} f(z) = w_1$

Then $|f(z) - w_0| < \varepsilon$ whenever $0 < |z - z_0| < \delta_0$

$|f(z) - w_1| < \varepsilon$ whenever $0 < |z - z_0| < \delta_1$

Let $\delta = \min(\delta_0, \delta_1)$

if $0 < |z - z_0| < \delta$

$$|[f(z) - w_0] - [f(z) - w_1]| \leq |f(z) - w_0| + |f(z) - w_1| < 2\varepsilon$$

$$|w_1 - w_0| < 2\varepsilon$$

Hence $|w_1 - w_0|$ is a nonnegative constant, and can be chosen arbitrarily small.


$$w_1 - w_0 = 0, \quad \text{or} \quad w_1 = w_0$$

Ex 2. If $f(z) = \frac{z}{z}$ (4)
 then $\lim_{z \rightarrow 0} f(z)$ does not exist.

$$\lim_{z \rightarrow 0} f(z)$$

show: when $z = (x, 0)$ $f(z) = \frac{x + i0}{x - i0} = 1$

when $z = (0, y)$ $f(z) = \frac{0 + iy}{0 - iy} = -1$



since a limit is unique, limit of (4) does not exist.

(2) provides a means of testing whether a given point W_0 is a limit, it does not directly provide a method for determining that limit.

Theorems on limits

Thm 1. Suppose that

$$f(z) = u(x, y) + iv(x, y), \quad z_0 = x_0 + iy_0$$

and $w_0 = u_0 + iv_0$

Then $\lim_{z \rightarrow z_0} f(z) = w_0$ iff

$$\lim_{(x, y) \rightarrow (x_0, y_0)} u(x, y) = u_0 \quad \text{and} \quad \lim_{(x, y) \rightarrow (x_0, y_0)} v(x, y) = v_0$$

pf: " \Leftarrow $|u - u_0| < \frac{\varepsilon}{2}$ whenever $0 < \sqrt{(x - x_0)^2 + (y - y_0)^2} < \delta_1$

$|v - v_0| < \frac{\varepsilon}{2}$ whenever $0 < \sqrt{(x - x_0)^2 + (y - y_0)^2} < \delta_2$

let $\delta = \min(\delta_1, \delta_2)$

since

$$|(u + iv) - (u_0 + iv_0)| = |(u - u_0) + i(v - v_0)| \leq |u - u_0| + |v - v_0|$$

and

$$\sqrt{(x - x_0)^2 + (y - y_0)^2} = |(x - x_0) + i(y - y_0)| = |(x + iy) - (x_0 + iy_0)|$$

$$\therefore |(u + iv) - (u_0 + iv_0)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon$$

$$\text{whenever } 0 < |(x + iy) - (x_0 + iy_0)| < \delta$$

“ ”

\Rightarrow

$$\text{But } |(u + iv) - (u_0 + iv_0)| < \varepsilon \quad \text{whenever } 0 < |(x + iy) - (x_0 + iy_0)| < \delta$$

$$|u - u_0| \leq |(u - u_0) + i(v - v_0)| = |(u + iv) - (u_0 + iv_0)| < \varepsilon$$

$$\text{and } |v - v_0| \leq |(u - u_0) + i(v - v_0)| = |(u + iv) - (u_0 + iv_0)| < \varepsilon$$

$$|(x + iy) - (x_0 + iy_0)| = \sqrt{(x - x_0)^2 + (y - y_0)^2}$$

$$\therefore |u - u_0| < \varepsilon \quad \text{and} \quad |v - v_0| < \varepsilon$$

$$\text{whenever } 0 < \sqrt{(x - x_0)^2 + (y - y_0)^2} < \delta$$

Thm 2. suppose that

$$\lim_{z \rightarrow z_0} f(z) = w_0 \quad \text{and} \quad \lim_{z \rightarrow z_0} F(z) = W_0 \quad (7)$$

Then $\lim_{z \rightarrow z_0} [f(z) + F(z)] = w_0 + W_0$

$$\lim_{z \rightarrow z_0} [f(z) \cdot F(z)] = w_0 W_0 \quad (9)$$

and if $W_0 \neq 0$

$$\lim_{z \rightarrow z_0} \frac{f(z)}{F(z)} = \frac{w_0}{W_0}$$

pf: utilize Thm 1.

for (9). $f(z) = u(x, y) + iv(x, y)$

$$F(z) = U(x, y) + iV(x, y)$$

$$z_0 = x_0 + iy_0, \quad w_0 = u_0 + iv_0, \quad W_0 = U_0 + iV_0$$

use Thm 1. and (7)

$f(z)F(z) = (uU - vV) + i(vU + uV)$ have the limits

$$\begin{array}{ccc} & \Downarrow & \Downarrow \\ & u_0U_0 - v_0V_0 & v_0U_0 + u_0V_0 \\ & = w_0W_0 & \end{array}$$

An immediate consequence of Thm. 1:

• $\lim_{z \rightarrow z_0} c = c$

• $\lim_{z \rightarrow z_0} z = z_0$

• $\lim_{z \rightarrow z_0} z^n = z_0^n \quad (n = 1, 2, \dots)$

by property (9) and math induction.

• $P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n \quad (11)$

$\lim_{z \rightarrow z_0} P(z) = P(z_0)$


• if $\lim_{z \rightarrow z_0} f(z) = w_0$, then $\lim_{z \rightarrow z_0} |f(z)| = |w_0|$

$\left| |f(z)| - |w_0| \right| \leq |f(z) - w_0| < \varepsilon$ whenever $0 < |z - z_0| < \delta$

Limits involving the point at Infinity

It is sometime convenient to include with the complex plane the point at infinity, denoted by ∞ , and to use limits involving it.

Complex plane + infinity = extended complex plane.



complex plane passing thru the equator of a unit sphere.

To each point z in the plane there corresponds exactly one point P on the surface of the sphere.




intersection of the line z - N with the surface.

↑
north pole

To each point P on the surface of the sphere, other than the north pole N , there corresponds exactly one point z in the plane.

By letting the point N of the sphere correspond to the point at infinity, we obtain a one-to-one correspondence between the points of the sphere and the points of the extended complex plane.



- $\lim_{z \rightarrow z_0} f(z) = \infty$

$$\Leftrightarrow |f(z)| > \frac{1}{\varepsilon} \quad \text{whenever} \quad 0 < |z - z_0| < \delta$$

$$\Leftrightarrow \left| \frac{1}{f(z)} - 0 \right| < \varepsilon \quad \text{whenever} \quad 0 < |z - z_0| < \delta$$

$$\therefore \lim_{z \rightarrow z_0} f(z) = \infty \quad \text{iff} \quad \lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0$$

Ex1. $\lim_{z \rightarrow -1} \frac{iz + 3}{z + 1} = \infty$ since $\lim_{z \rightarrow -1} \frac{z + 1}{iz + 3} = 0$

- $\lim_{z \rightarrow \infty} f(z) = w_0$

$$\Leftrightarrow |f(z) - w_0| < \varepsilon \quad \text{whenever} \quad |z| > \frac{1}{\delta}$$

$$\Leftrightarrow \left| f\left(\frac{1}{z}\right) - w_0 \right| < \varepsilon \quad \text{whenever} \quad 0 < |z - 0| < \delta$$

$$\therefore \lim_{z \rightarrow \infty} f(z) = w_0 \quad \text{iff} \quad \lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) = w_0$$

EX 2. $\lim_{z \rightarrow \infty} \frac{2z + i}{z + 1} = 2$ since $\lim_{z \rightarrow 0} \frac{\frac{2}{z} + i}{\frac{1}{z} + 1} = \lim_{z \rightarrow 0} \frac{2 + iz}{1 + z} = 2$

- $\lim_{z \rightarrow \infty} f(z) = \infty$

$$\Leftrightarrow |f(z)| > \frac{1}{\varepsilon} \quad \text{whenever} \quad |z| > \frac{1}{\delta}$$

$$\Leftrightarrow \left| f\left(\frac{1}{z}\right) \right| > \frac{1}{\varepsilon} \quad \text{whenever} \quad \left| \frac{1}{z} \right| > \frac{1}{\delta}$$

$$\Leftrightarrow \left| \frac{1}{f(1/z)} - 0 \right| < \varepsilon \quad \text{whenever} \quad 0 < |z - 0| < \delta$$

$$\therefore \lim_{z \rightarrow \infty} f(z) = \infty \quad \text{iff} \quad \lim_{z \rightarrow 0} \frac{1}{f\left(\frac{1}{z}\right)} = 0$$

Ex 3. $\lim_{z \rightarrow \infty} \frac{2z^3 - 1}{z^2 + 1} = \infty$

since $\lim_{z \rightarrow 0} \frac{\frac{1}{z^2} + 1}{\frac{2}{z^3} - 1} = \lim_{z \rightarrow 0} \frac{z + z^3}{2 - z^3} = 0$

Continuity

A function f is continuous at a point z_0 if

$$\lim_{z \rightarrow z_0} f(z) \text{ exists,} \quad (1)$$

$$f(z_0) \text{ exists,} \quad (2)$$


$$\lim_{z \rightarrow z_0} f(z) = f(z_0) \quad (3) \quad ((3) \text{ implies } (1)(2))$$

$$(|f(z) - f(z_0)| < \varepsilon \quad \textit{whenever} \quad |z - z_0| < \delta)$$

- if f_1, f_2 continuous at z_0 , then $f_1 + f_2, f_1 f_2$
also continuous at z_0 .

$$\text{So is } \frac{f_1}{f_2} \quad \textit{if} \quad f_2(z_0) \neq 0$$

- A polynomial is continuous in the entire plane because of (11), section 12. p.37
- A composition of continuous function is continuous.



$$|g[f(z)] - g[f(z_0)]| < \varepsilon, \quad \text{whenever } |f(z) - f(z_0)| < r,$$

$$\text{whenever } |z - z_0| < \delta$$

- If a function $f(z)$ is continuous and non zero at a point z_0 , then $f(z) \neq 0$ throughout some neighborhood of that point.

when $f(z_0) \neq 0$ let $\varepsilon = \frac{|f(z_0)|}{2}$

$$|f(z) - f(z_0)| < \frac{|f(z_0)|}{2} \quad \text{whenever } |z - z_0| < \delta$$

if there is a point z in the $|z - z_0| < \delta$ at which $f(z) = 0$

$$|f(z_0)| < \frac{|f(z_0)|}{2} \quad \text{a contradiction.}$$

From Thm 1

a function f of a complex variable is continuous at a point
iff its component functions u and v are continuous there.

$$z_0 = (x_0, y_0)$$

Ex. The function

$f(z) = \cos(x^2 - y^2) \cosh 2xy - i \sin(x^2 - y^2) \sinh 2xy$
is continuous everywhere in the complex plane since

- (i) $x^2 - y^2$ are continuous (polynomial)
 $2xy$
- (ii) \cos, \sin, \cosh, \sinh are continuous
- (iii) real and imaginary component are continuous

complex function is continuous.



$$\sinh x = \frac{e^x - e^{-x}}{2} \quad \sin x = \frac{e^{ix} - e^{-ix}}{2i}$$
$$\cosh x = \frac{e^x + e^{-x}}{2} \quad \cos x = \frac{e^{ix} + e^{-ix}}{2}$$

Derivatives

Let f be a function whose domain of definition $f'(z_0)$ contain a neighborhood of a point z_0 . The derivative of f at z_0 , written $f'(z_0)$, is

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

provided this limit exists.


f is said to be differentiable at z_0 .

let $\Delta z = z - z_0$

$$f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$$

let $\Delta w = f(z + \Delta z) - f(z)$.

$$f'(z) = \frac{dw}{dz} = \lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z}$$



Ex1. Suppose $f(z) = z^2$ at any point z

$$\lim_{\Delta z \rightarrow 0} \frac{\Delta w}{\Delta z} = \lim_{\Delta z \rightarrow 0} \frac{(z + \Delta z)^2 - z^2}{\Delta z} = \lim_{\Delta z \rightarrow 0} (2z + \Delta z) = 2z + 0 = 2z$$

since $2z + \Delta z$ a polynomial in Δz .

$$\therefore f'(z) = \frac{dw}{dz} = 2z$$

Ex2. $f(z) = |z|^2$


$$\frac{\Delta w}{\Delta z} = \frac{|z + \Delta z|^2 - |z|^2}{\Delta z} = \frac{(z + \Delta z)(\bar{z} + \overline{\Delta z}) - z\bar{z}}{\Delta z} = \bar{z} + \overline{\Delta z} + z \frac{\overline{\Delta z}}{\Delta z}$$

when $\Delta z \rightarrow 0$ thru $(\Delta x, 0)$ on the real axis $\overline{\Delta z} = \Delta z$

Hence if the limit of $\frac{\Delta w}{\Delta z}$ exists, its value = $\bar{z} + z$

when $\Delta z \rightarrow 0$ thru $(0, \Delta y)$ on the imaginary axis.

$\overline{\Delta z} = -\Delta z$, limit = $\bar{z} - z$ if it exists.



since limits are unique,

$$\overline{z} + z = \overline{z} - z, \quad \text{or} \quad z = 0 \quad \text{if} \quad \frac{dw}{dz} \text{ is to exist.}$$

observe that $\frac{\Delta w}{\Delta z} \rightarrow \overline{\Delta z}$ when $z = 0$

$$\therefore \frac{dw}{dz} \text{ exists only at } z = 0, \text{ its value} = 0$$

- Example 2 shows that

a function can be differentiable at a certain point but nowhere else in any neighborhood of that point.


- Re $|z|^2 = x^2 + y^2$ are continuous, partially

Im $|z|^2 = 0$ differentiable at a point.


but $|z|^2$ may not be differentiable there.

- $f(z) = |z|^2$ is continuous at each point in the plane since its components are continuous at each point.

∴ continuity $\xrightarrow{\text{not necessarily}}$ derivative exists.



existence of derivative \Rightarrow continuity.



$$\begin{aligned} \lim_{z \rightarrow z_0} [f(z) - f(z_0)] &= \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \lim_{z \rightarrow z_0} (z - z_0) \\ &= f'(z_0) \cdot 0 = 0 \\ \therefore \lim_{z \rightarrow z_0} f(z) &= f(z_0) \end{aligned}$$

16. Differentiation Formulas

$$\frac{d}{dz} C = 0 \quad C : \text{complex constant}$$

$$\frac{d}{dz} z = 1$$

$$\frac{d}{dz} [cf(z)] = cf'(z)$$

$$\frac{d}{dz} z^n = nz^{n-1} \quad n \text{ a positive integer.}$$

$$\frac{d}{dz} [f(z) + F(z)] = f'(z) + F'(z)$$

$$\frac{d}{dz} [f(z)F(z)] = f(z)F'(z) + f'(z)F(z) \quad (4)$$

when $F(z) \neq 0$

$$\frac{d}{dz} \left[\frac{f(z)}{F(z)} \right] = \frac{F(z)f'(z) - f(z)F'(z)}{[F(z)]^2}$$

pf : (4)

$$f(z + \Delta z)F(z + \Delta z) - f(z)F(z)$$

$$= f(z)[F(z + \Delta z) - F(z)] + [f(z + \Delta z) - f(z)]F(z + \Delta z)$$

$$\frac{f(z + \Delta z)F(z + \Delta z) - f(z)F(z)}{\Delta z} = f(z)\frac{F(z + \Delta z) - F(z)}{\Delta z} + \frac{f(z + \Delta z) - f(z)}{\Delta z}F(z + \Delta z)$$

$$\text{as } \Delta z \rightarrow 0 \quad \frac{d}{dz}[fF] = f(z)F'(z) + f'(z)F(z + \Delta z)$$

$$= f(z)F'(z) + f'(z)F(z) \quad (F \text{ continuous at } z)$$

f has a derivative at z_0

g has a derivative at $f(z_0)$

$F(z)=g[f(z)]$ has a derivative at z_0

and $F'(z_0) = g'[f(z_0)]f'(z_0)$ chain rule (6)

$$\frac{dW}{dz} = \frac{dW}{dw} \frac{dw}{dz}$$

pf of (6)

choose a z_0 at which $f'(z_0)$ exists.

let $w_0 = f(z_0)$ and assume $g'(w_0)$ exists.

Then, there is $|w - w_0| < \varepsilon$ of w_0 such that

we can define a function Φ , with $\Phi(w_0) = 0$

$$\Phi(w) = \frac{g(w) - g(w_0)}{w - w_0} - g'(w_0) \quad \text{when } w \neq w_0 \quad (7)$$

$\lim_{w \rightarrow w_0} \Phi(w) = 0$, Hence Φ is continuous at w_0

$$(7) \Rightarrow \quad g(w) - g(w_0) = [g'(w_0) + \Phi(w)](w - w_0) \quad (|w - w_0| < \varepsilon) \quad (9)$$

valid even when $w = w_0$

since $f'(z_0)$ exists and therefore f is continuous at z_0 , then we can

have $f(z)$ lies in $|w - w_0| < \varepsilon$ of w_0 if $|z - z_0| < \delta$

substitute w by $f(z)$ in (9) when z in $|z - z_0| < \delta$

(9) becomes

$$\frac{g[f(z)] - g[f(z_0)]}{z - z_0} = \{g'[f(z_0)] + \Phi[f(z)]\} \frac{f(z) - f(z_0)}{z - z_0} \quad (10)$$

$$(0 < |z - z_0| < \delta)$$

since f is continuous at z_0 , Φ is continuous at $w_0 = f(z_0)$

$\therefore \Phi[f(z)]$ is continuous at z_0 , and since $\Phi(w_0) = 0$

$$\lim_{z \rightarrow z_0} \Phi[f(z)] = 0$$

so (10) becomes $F'(z_0) = g'[f(z_0)] f'(z_0)$ as $z \rightarrow z_0$

Cauchy-Riemann Equations

Suppose that $f'(z_0) = \lim_{\Delta z \rightarrow 0} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}$ exists.

writing $z_0 = x_0 + iy_0$, $\Delta z = \Delta x + i\Delta y$

Then by Thm. 1

$$\operatorname{Re}[f'(z_0)] = \lim_{(\Delta x, \Delta y) \rightarrow (0,0)} \operatorname{Re}\left[\frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}\right] \quad (3)$$

$$\operatorname{Im}[f'(z_0)] = \lim_{(\Delta x, \Delta y) \rightarrow (0,0)} \operatorname{Im}\left[\frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z}\right] \quad (4)$$

where

$$\begin{aligned} \frac{f(z_0 + \Delta z) - f(z_0)}{\Delta z} &= \frac{u(x_0 + \Delta x, y_0 + \Delta y) - u(x_0, y_0)}{\Delta x + i\Delta y} \\ &\quad + \frac{i[v(x_0 + \Delta x, y_0 + \Delta y) - v(x_0, y_0)]}{\Delta x + i\Delta y} \end{aligned} \quad (5)$$

Let $(\Delta x, \Delta y)$ tend to $(0,0)$ horizontally through $(\Delta x, 0)$ $\Delta y = 0$

$$\therefore \operatorname{Re}[f'(z_0)] = \lim_{\Delta x \rightarrow 0} \frac{u(x_0 + \Delta x, y_0) - u(x_0, y_0)}{\Delta x}$$

$$\operatorname{Im}[f'(z_0)] = \lim_{\Delta x \rightarrow 0} \frac{v(x_0 + \Delta x, y_0) - v(x_0, y_0)}{\Delta x}$$

$$\therefore f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0) \quad (6)$$

Let $(\Delta x, \Delta y)$ tend to $(0,0)$ vertically through $(0, \Delta y)$ $\Delta x = 0$

$$f'(z_0) = \left(\frac{u(x_0, y_0 + \Delta y) - u(x_0, y_0)}{i\Delta y} + \frac{i[v(x_0, y_0 + \Delta y) - v(x_0, y_0)]}{i\Delta y} \right)$$

$$= v_y(x_0, y_0) - iu_y(x_0, y_0) \quad (7)$$

$$= -iu_y + v_y$$

$$(6) = (7)$$

$$\therefore u_x(x_0, y_0) = v_y(x_0, y_0) \quad (8)$$

$$u_y(x_0, y_0) = -v_x(x_0, y_0)$$

Cauchy-Riemann Equations.

Thm : suppose $f(z) = u(x, y) + iv(x, y)$

$f'(z)$ exists at a point $z_0 = x_0 + iy_0$

Then u_x, u_y, v_x, v_y exist at (x_0, y_0)

and $u_x = v_y, u_y = -v_x$; also $f'(z) = u_x + iv_x$

Ex 1. $f(z) = z^2 = x^2 - y^2 + i2xy$

$$u_x = 2x \quad v_x = 2y$$

$$u_y = -2y \quad v_y = 2x$$

$$u_x = v_y, \quad u_y = -v_x$$

$$f'(z) = 2x + i2y = 2(x + iy) = 2z$$

Cauchy-Riemann equations are Necessary conditions for the existence of the derivative of a function f at z_0 .

➡ Can be used to locate points at which f does not have a derivative.

Ex 2. $f(z) = |z|^2$,

$$u(x, y) = x^2 + y^2 \quad v(x, y) = 0$$

$$u_x = 2x \quad v_x = 0 \quad u_x \neq v_y, \quad f'(z) \text{ does not exist}$$

$$u_y = 2y \quad v_y = 0 \quad \text{at any nonzero point.}$$

The above Thm does not ensure the existence of $f'(z_0)$
(say)

Sufficient Conditions For Differentiability

$$f'(z_0) \text{ exist} \rightarrow u_x = v_y, \quad u_y = -v_x$$

but not
"←"

Thm.

Let $f(z) = u(x, y) + iv(x, y)$ be defined throughout some neighborhood of a point

$$z_0 = x_0 + iy_0$$

suppose u_x, u_y, v_x, v_y exist everywhere in the neighborhood and are **continuous** at (x_0, y_0)

Then, if $u_x = v_y, \quad u_y = -v_x$ at (x_0, y_0)
 $\Rightarrow f'(z_0)$ exists.

pf : let $\Delta z = \Delta x + i\Delta y$, where $0 < |\Delta z| < \varepsilon$

$$\Delta w = f(z_0 + \Delta z) - f(z_0) \quad \curvearrowright$$

Thus $\Delta w = \Delta u + i\Delta v \iff u(z_0 + \Delta z) - u(z_0) + i[v(z_0 + \Delta z) - v(z_0)]$

where $\Delta u = u(x_0 + \Delta x, y_0 + \Delta y) - u(x_0, y_0)$

$$\Delta v = v(x_0 + \Delta x, y_0 + \Delta y) - v(x_0, y_0)$$

⇒ Now in view of the continuity of the first-order partial derivatives of u and v at the point (x_0, y_0)

$$\begin{aligned} \Delta u = & u(x_0, y_0) + u_x(x_0, y_0)\Delta x + u_y(x_0, y_0)\Delta y + u_{xy}(x_0, y_0)\Delta x\Delta y \\ & + u_{xx}(x_0, y_0)\frac{\Delta x^2}{2!} \\ & + u_{yy}(x_0, y_0)\frac{\Delta y^2}{2!} \\ & - u(x_0, y_0) \qquad \qquad \qquad + \dots \end{aligned}$$

$$= u_x(x_0, y_0)\Delta x + u_y(x_0, y_0)\Delta y + \varepsilon_1 \sqrt{(\Delta x)^2 + (\Delta y)^2}$$

$$\Delta v = v_x(x_0, y_0)\Delta x + v_y(x_0, y_0)\Delta y + \varepsilon_2 \sqrt{(\Delta x)^2 + (\Delta y)^2}$$

$$\varepsilon_1, \varepsilon_2 \rightarrow 0, \text{ as } (\Delta x, \Delta y) \rightarrow (0, 0)$$

$$\Delta w = \Delta u + i\Delta v \quad \leftarrow \text{ where } \varepsilon_1 \text{ and } \varepsilon_2 \text{ tend to 0 as } (\Delta x, \Delta y) \rightarrow (0, 0) \text{ in the } z\text{-plane.}$$

$$= \text{above} \quad (3)$$

assuming that the Cauchy-Riemann equations are satisfied at (x_0, y_0) we can replace

$$u_y \text{ by } -v_x, \text{ and } v_y \text{ by } u_x \quad (3) \quad \Delta z$$

to get

$$\frac{\Delta w}{\Delta z} = u_x(x_0, y_0) + iv_x(x_0, y_0) + (\varepsilon_1 + i\varepsilon_2) \frac{\sqrt{(\Delta x)^2 + (\Delta y)^2}}{\Delta z} \quad (4)$$

$$\text{but } \sqrt{(\Delta x)^2 + (\Delta y)^2} = |\Delta z|$$

$$\text{so } \left| \frac{\sqrt{(\Delta x)^2 + (\Delta y)^2}}{\Delta z} \right| = 1$$

also $\varepsilon_1 + i\varepsilon_2$ tends to 0, as $(\Delta x, \Delta y) \rightarrow (0, 0)$

The last term in (4) tends to 0 as $\Delta z \rightarrow 0$

\therefore The limit of $\frac{\Delta w}{\Delta z}$ exists, and $f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0)$.

Ex 1. $f(z) = e^x (\cos y + i \sin y)$

$$u(x, y) = e^x \cos y$$

$$v(x, y) = e^x \sin y$$

$$u_x = v_y, \quad u_y = -v_x \quad \text{everywhere, and continuous.}$$

$\Rightarrow f'(z)$ exists everywhere, and

$$f'(z) = u_x + iv_x = e^x (\cos y + i \sin y)$$

Ex 2. $f(z) = |z|^2$

$$u(x, y) = x^2 + y^2 \quad u_x = 2x \quad u_y = 2y$$

$$v(x, y) = 0 \quad v_x = 0 \quad v_y = 0$$

has a derivative at $z=0$.

$$f'(0) = 0 + i0$$

can not have derivative at any nonzero point.

Polar Coordinates

$$x = r \cos \theta \quad y = r \sin \theta$$

$$z = x + iy = re^{i\theta} \quad (z \neq 0)$$

Suppose that u_x, u_y, v_x, v_y exist everywhere in some neighborhood of a given non-zero point z_0 and are continuous at that point.

$u_r, u_\theta, v_r, v_\theta$ also have these properties, and (by chain rule)

$$\frac{\partial u}{\partial r} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial r}$$

$$\frac{\partial u}{\partial \theta} = \frac{\partial u}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial u}{\partial y} \frac{\partial y}{\partial \theta}$$

$$u_r = u_x \cos \theta + u_y \sin \theta \quad (2)$$

$$u_\theta = -u_x r \sin \theta + u_y r \cos \theta$$

Similarly,

$$v_r = v_x \cos \theta + v_y \sin \theta \quad (3)$$

$$v_\theta = -v_x r \sin \theta + v_y r \cos \theta$$

If $u_x = v_y, u_y = -v_x$

$$v_r = -u_y \cos \theta + u_x \sin \theta \quad (5)$$

$$v_\theta = u_y r \sin \theta + u_x r \cos \theta$$

from (2) (5), $u_r = \frac{1}{r} v_\theta$ at z_0 (6)

$$\frac{1}{r} u_\theta = -v_r$$

Thm. p53...

$$f'(z_0) = u_x + iv_x$$

$$= ?$$

$$u_r = u_x \cos \theta + u_y \sin \theta$$

$$u_r \cos \theta = u_x \cos^2 \theta + u_y \sin \theta \cos \theta$$

$$\boxed{\therefore u_r \cos \theta + v_r \sin \theta = u_x}$$

$$u_r = v_y \cos \theta - v_x \sin \theta$$

$$u_r \sin \theta = v_y \cos \theta \sin \theta - v_x \sin^2 \theta$$

$$\boxed{v_r \cos \theta - u_r \sin \theta = v_x}$$

$$v_r = v_x \cos \theta + v_y \sin \theta$$

$$= -u_y \cos \theta + u_x \sin \theta$$

$$v_r \sin \theta = -u_y \cos \theta \sin \theta + u_x \sin^2 \theta$$

$$v_r = v_x \cos \theta + v_y \sin \theta$$

$$\cos \theta v_r = v_x \cos^2 \theta + v_y \sin \theta \cos \theta$$

$$\begin{aligned} \therefore f'(z_0) &= u_r \cos \theta + v_r \sin \theta + i(v_r \cos \theta - u_r \sin \theta) \\ &= (\cos \theta - i \sin \theta)(u_r + iv_r) \\ &= e^{-i\theta} (u_r + iv_r) \end{aligned} \quad (7)$$

Ex : Consider $f(z) = \frac{1}{z} = \frac{1}{re^{i\theta}}$

$$u(r, \theta) = \frac{1}{r} \cos \theta \quad v(r, \theta) = -\frac{1}{r} \sin \theta$$

$$u_r = -\frac{1}{r^2} \cos \theta \quad v_r = \frac{1}{r^2} \sin \theta$$

$$u_\theta = -\frac{1}{r} \sin \theta \quad v_\theta = -\frac{1}{r} \cos \theta$$

$$\Rightarrow u_r = \frac{1}{r} v_\theta, \quad \frac{1}{r} u_\theta = -v_r \quad \text{at any non-zero point} \quad z = re^{i\theta}$$

$\therefore f'$ exists

$$f' = e^{-i\theta} \left(-\frac{1}{r^2} \cos \theta + \frac{i}{r^2} \sin \theta \right)$$

$$= \frac{1}{r^2} (-e^{-i\theta}) e^{-i\theta} = -\frac{1}{r^2} e^{-i2\theta} = -\frac{1}{z^2}$$