

# Expansion of Functions of Several Variables

## Functions of Several Variables

We have studied functions of one variable,  $y = f(x)$  in which  $x$  was the independent variable and  $y$  was the dependent variable. We are going to expand the idea of functions to include functions with more than one independent variable. For example, consider the functions below:

$$f(x, y) = 2x^2 + y^2$$

or

$$g(x, y, z) = 2xe^{yz}$$

or

$$h(x_1, x_2, x_3, x_4) = 2x_1 - x_2 + 4x_3 + x_4$$

Hopefully you can see the notation for functions of several variables is similar to the notation you've used with single variable functions.

$$z = f(x, y) = 2x^2 + y^2$$

or

$$w = g(x, y, z) = 2xe^{yz}$$

or

$$h(x_1, x_2, x_3, x_4) = 2x_1 - x_2 + 4x_3 + x_4$$

The function  $z = f(x, y)$  is a function of two variables. It has independent variables  $x$  and  $y$ , and the dependent variable  $z$ .

Likewise, the function  $w = g(x, y, z)$  is a function of three variables. The variables  $x$ ,  $y$  and  $z$  are independent variables and  $w$  is the dependent variable.

The function  $h$  is similar except there are four independent variables.

When finding values of the several variable functions instead of just substituting in an x-value, we will substitute in values for each of the independent variables:

For example, using the function  $f$  on the previous slide, we will evaluate the function  $f(x, y)$  for  $(2, 3)$ ,  $(4, -3)$  and  $(5, y)$ .

$$f(x, y) = 2x^2 + y^2$$

$$f(2, 3) = 2 \cdot 2^2 + 3^2 = 2 \cdot 4 + 9 = 17$$

$$f(4, -3) = 2 \cdot 4^2 + (-3)^2 = 2 \cdot 16 + 9 = 41$$

$$f(5, y) = 2 \cdot 5^2 + y^2 = 2 \cdot 25 + y^2 = 50 + y^2$$

A Function of Two Variables: A function  $f$  of two variables  $x$  and  $y$  is a rule that assigns to each ordered pair  $(x, y)$  in a given set  $D$ , called the domain, a unique value of  $f$ .

Functions of more variables can be defined similarly.

The operations we performed with one-variable functions can also be performed with functions of several variables.

For example, for the two-variable functions  $f$  and  $g$ :

$$(f \pm g)(x, y) = f(x, y) \pm g(x, y)$$

$$(f \cdot g)(x, y) = f(x, y) \cdot g(x, y)$$

$$\left(\frac{f}{g}\right)(x, y) = \frac{f(x, y)}{g(x, y)}, \text{ Provided } g(x, y) \neq 0$$

In general we will not consider the composition of two multi-variable functions.

## Domains of Functions of Several Variables:

Unless the domain is given, assume the domain is the set of all points for which the equation is defined.

For example, consider the functions

$$f(x,y) = 3x^2 + y^2 \quad \text{and} \quad g(x,y) = \frac{1}{\sqrt{xy}}$$

The domain of  $f(x,y)$  is the entire  $xy$ -plane. Every ordered pair in the  $xy$ -plane will produce a real value for  $f$ .


The domain of  $g(x,y)$  is the set of all points  $(x,y)$  in the  $xy$ -plane such that the product  $xy$  is greater than 0. This would be all the points in the first quadrant and the third quadrant.

Example 1: Find the domain of the function:  $f(x, y) = \sqrt{25 - x^2 - y^2}$

Solution: The domain of  $f(x, y)$  is the set of all points that satisfy the inequality:

$$25 - x^2 - y^2 \geq 0 \quad \text{or} \quad 25 \geq x^2 + y^2$$

You may recognize that this is similar to the equation of a circle and the inequality implies that any ordered pair on the circle or inside the circle  $x^2 + y^2 = 25$  is in the domain.



The highlighted area is the domain to  $f$ .

Example 2: Find the domain of the function:

$$g(x, y, z) = \sqrt{x^2 + y^2 + z^2 - 16}$$

Solution:

Note that  $g$  is a function of three variables, so the domain is NOT an area in the  $xy$ -plane. The domain of  $g$  is a solid in the 3-dimensional coordinate system.

The expression under the radical must be nonnegative, resulting in the inequality:

$$x^2 + y^2 + z^2 - 16 \geq 0 \text{ or } x^2 + y^2 + z^2 \geq 16$$

This implies that any ordered triple outside of the sphere centered at the origin with radius 4 is in the domain.




Example 3: Find the domain of the function:  $h(x,y) = \ln(xy)$

Solution:

We know the argument of the natural log must be greater than zero.

$$\text{So, } x \cdot y > 0$$

This occurs in quadrant I and quadrant III. The domain is highlighted below. Note the x-axis and the y-axis are NOT in the domain.



## Graphs of Functions of Several Variables

As you learned in 2-dimensional space the graph of a function can be helpful to your understanding of the function. The graph gives an illustration or visual representation of all the solutions to the equation. We also want to use this tool with functions of two variables.

The graph of a function of two variables,  $z = f(x, y)$ , is the set of ordered triples,  $(x, y, z)$  for which the ordered pair,  $(x, y)$  is in the domain.

\*The graph of  $z = f(x, y)$  is a surface in 3-dimensional space.

The graph of a function of three variables,  $w = f(x, y, z)$  is the set of all points  $(x, y, z, w)$  for which the ordered triple,  $(x, y, z)$  is in the domain.

\*The graph of  $w = f(x, y, z)$  is in 4 dimensions.

We can't draw this graph or the graphs of any functions with 3 or more independent variables.

Example 4: Find the domain and range of the function and then sketch the graph.

$$z = f(x, y) = \sqrt{25 - x^2 - y^2}$$

Solution: From Example 1 we know the domain is all ordered pairs  $(x, y)$  on or inside the circle centered at the origin with radius 5.

All ordered pairs satisfying the inequality:  $x^2 + y^2 \leq 25$

The range is going to consist of all possible outcomes for  $z$ .

The range must be nonnegative since  $z$  equals a principle square root and furthermore, with the domain restriction:  $x^2 + y^2 \leq 25$ , the value of the radicand will only vary between 0 and 25.

Thus, the range is  $0 \leq z \leq 5$ .

### Solution to Example 4 Continued:

Now let's consider the sketch of the function:  $z = \sqrt{25 - x^2 - y^2}$

Squaring both sides and simplifying:  $z^2 = 25 - x^2 - y^2$

$$x^2 + y^2 + z^2 = 25$$

You may recognize this equation from Chapter 7 - A sphere with radius 5. This is helpful to sketching the function, but we must be careful !!

The **function**  $z = \sqrt{25 - x^2 - y^2}$  and the **equation**  $x^2 + y^2 + z^2 = 25$

are not exactly the same. The **equation** does NOT represent  $z$  as a function of  $x$  and  $y$  - meaning there is not a unique value for  $z$  for each  $(x, y)$ . Keep in mind that the **function** had a range of  $0 \leq z \leq 5$ , which means the function is only the top half of the sphere.

As you have done before when sketching a surface in 3-dimensions it may be helpful for you to use the traces in each coordinate plane.

1. The trace in the  $xy$ -plane,  $z = 0$ , is the equation:

$$0 = \sqrt{25 - x^2 - y^2} \quad \text{or} \quad x^2 + y^2 = 25$$

The circle centered at the origin with radius 5 in the  $xy$ -plane.

2. The trace in the  $yz$ -plane,  $x = 0$ , is the equation:

$$z = \sqrt{25 - y^2} \quad \text{or} \quad y^2 + z^2 = 25$$

The circle centered at the origin with radius 5 in the  $yz$ -plane.

3. The trace in the  $xz$ -plane,  $y = 0$ , is the equation:

$$z = \sqrt{25 - x^2} \quad \text{or} \quad x^2 + z^2 = 25$$

The circle centered at the origin with radius 5 in the  $xz$ -plane.

Along with sketching the traces in each coordinate plane, it may be helpful to sketch traces in planes parallel to the coordinate planes.

4. Let  $z = 3$ :  $3 = \sqrt{25 - x^2 - y^2}$  or  $x^2 + y^2 = 16$

So on the plane  $z = 3$ , parallel to the  $xy$ -plane, the trace is a circle centered at  $(0,0,3)$  with radius 4.

5. Let  $z = 4$ :  $4 = \sqrt{25 - x^2 - y^2}$  or  $x^2 + y^2 = 9$

So on the plane  $z = 4$ , parallel to the  $xy$ -plane, the trace is a circle centered at  $(0,0,4)$  with radius 3.