

Inverse of a Matrix



The inverse is Defined as $AA^{-1} = A^{-1}A = I$

$$\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \begin{bmatrix} ? & ? \\ ? & ? \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} \quad \begin{bmatrix} ? & ? \\ ? & ? \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$$

We will find out how to calculate the inverse for 2x2 matrix

But first why is it important ?

Because it will allow us to solve equations of the form

$$\begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \cdot \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$$

We will only consider 2x2
Matrix systems
That means simultaneous
equations

Because if we can express a system of equations in the form

$$Ax = b$$

Then we can multiply both sides by the inverse matrix

$$A^{-1}Ax = A^{-1}b$$

And we can then know the values of x because

$$A^{-1}A = I$$


$$x = A^{-1}b$$

$$\begin{pmatrix} 4 & -3 & 2 \\ -6 & 0 & 1 \end{pmatrix} \begin{matrix} \longrightarrow \\ \downarrow \end{matrix} \begin{pmatrix} 2 \\ -1 \\ 6 \end{pmatrix} = \begin{pmatrix} 8+3+12 \\ -12+0+6 \end{pmatrix} = \begin{pmatrix} 23 \\ -6 \end{pmatrix}$$

$$\begin{pmatrix} 6 & 3 \\ 1 & 2 \\ 5 & 7 \end{pmatrix} \begin{pmatrix} 2 \\ 3 \end{pmatrix} = \begin{pmatrix} 12+9 \\ 2+6 \\ 10+21 \end{pmatrix} = \begin{pmatrix} 21 \\ 8 \\ 31 \end{pmatrix}$$

$$\begin{pmatrix} 4 & 2 \\ 6 & 5 \\ 1 & 4 \end{pmatrix} = \begin{pmatrix} 24+10 & 4+8 \\ 34 & 12 \end{pmatrix}$$

* Dimensions and compatibility given by the domino rule



Note matrix multiplication is **not** commutative.

If A is a 3×1 and B is a 1×3 then AB is 3×3 BA is 1×1

- * This can only be done with SQUARE matrices
- * By hand we will only do this for a 2x2 matrix
- * Inverses of larger square matrices can be calculated but can be quite time expensive for large matrices, computers are generally used

$$\left(\begin{array}{c} \text{Ex} \\ \left(\begin{array}{cc} 4 & 8 \\ 1 & 3 \end{array} \right) \end{array} \right) A = \quad \text{then } A^{-1} = \left(\begin{array}{cc} 0.75 & -2 \\ -0.25 & 1 \end{array} \right) \quad \text{as } A \times A^{-1} = I$$

$$\left(\begin{array}{cc} 4 & 8 \\ 1 & 3 \end{array} \right) \left(\begin{array}{cc} 0.75 & -2 \\ -0.25 & 1 \end{array} \right) = \left(\begin{array}{cc} 3-2 & -8+8 \\ 0.75-0.75 & -2+3 \end{array} \right) = \left(\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array} \right)$$

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

Step-1 First find what is called the Determinant

This is calculated as $ad-bc$

Step-2 Then swap the elements in the leading diagonal

$$\begin{bmatrix} d & b \\ c & a \end{bmatrix}$$

Step-3 Then negate the other elements

$$\begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Step-4 Then multiply the Matrix by 1/determinant

$$\frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Step 1 - Calc Determinant

$$A = \begin{pmatrix} 4 & 8 \\ 1 & 3 \end{pmatrix} \quad \text{Determinant (ad-cb)} = 4 \times 3 - 8 \times 1 = 4$$

Step 2 - Swap Elements on leading diagonal

step2 $\begin{pmatrix} 3 & 8 \\ 1 & 4 \end{pmatrix}$

Step 3 - negate the other elements

step3 $\begin{pmatrix} 3 & -8 \\ -1 & 4 \end{pmatrix}$

Step 4 - multiply by 1/determinant

step4 $\frac{1}{4} \begin{pmatrix} 3 & -8 \\ -1 & 4 \end{pmatrix}$

$$A^{-1} = \begin{bmatrix} 0.75 & -2 \\ -0.25 & 1 \end{bmatrix}$$

check

$$\begin{aligned} AA^{-1} &= \begin{bmatrix} 4 & 8 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 0.75 & -2 \\ -0.25 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 3-2 & -8+8 \\ 0.75-0.75 & -2+3 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \end{aligned}$$

$$A = \begin{pmatrix} 2 & 6 \\ 1 & 5 \end{pmatrix} \quad A^{-1} = \frac{1}{4} \begin{pmatrix} 5 & -6 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 1.25 & -1.5 \\ -0.25 & 0.5 \end{pmatrix}$$

$$B = \begin{pmatrix} -5 & 20 \\ -1 & 2 \end{pmatrix} \quad B^{-1} = \frac{1}{10} \begin{pmatrix} 2 & -20 \\ 1 & -5 \end{pmatrix} = \begin{pmatrix} 0.2 & -2 \\ 0.1 & -0.5 \end{pmatrix}$$

$$C = \begin{pmatrix} 2 & 2 \\ 0 & -1 \end{pmatrix} \quad C^{-1} = \frac{1}{-2} \begin{pmatrix} -1 & -2 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 0.5 & 1 \\ 0 & -1 \end{pmatrix}$$

$$\mathbf{A} = \begin{pmatrix} 2 & 4 \\ 1 & 3 \end{pmatrix}$$

$$\mathbf{B} = \begin{pmatrix} 5 & 10 \\ -1 & 2 \end{pmatrix}$$

$$\mathbf{C} = \begin{pmatrix} 3 & 2 \\ 2 & 1 \end{pmatrix}$$

$$\mathbf{D} = \begin{pmatrix} 8 & 2 \\ -1 & 0 \end{pmatrix}$$

$$\mathbf{E} = \begin{pmatrix} 2 & -8 \\ -1 & 4 \end{pmatrix}$$

$$\mathbf{F} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\mathbf{A}^{-1} = \frac{1}{2} \begin{pmatrix} 3 & -4 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 1.5 & -2 \\ -0.5 & 1 \end{pmatrix}$$

$$\mathbf{B}^{-1} = \frac{1}{20} \begin{pmatrix} 2 & -10 \\ 1 & 5 \end{pmatrix} = \begin{pmatrix} 0.1 & -0.5 \\ 0.05 & 0.25 \end{pmatrix}$$

$$\mathbf{C}^{-1} = \frac{1}{-1} \begin{pmatrix} 1 & -2 \\ -2 & 3 \end{pmatrix} = \begin{pmatrix} -1 & 2 \\ 2 & -3 \end{pmatrix}$$

$$\mathbf{D}^{-1} = \frac{1}{2} \begin{pmatrix} 0 & -2 \\ 1 & 8 \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 0.5 & 4 \end{pmatrix}$$

E^{-1} cannot be found as $\det = 2 \times 4 - (-1 \times -8) = 8 - 8 = 0$

$$\mathbf{F}^{-1} = \frac{1}{1} \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \text{ self inverting}$$

- * Because matrices are clever storage systems for numbers there are a large and diverse number of ways we can apply them.
- * Matrices are used in to solve equations on computers
 - solving equations
- * They are used in computer games and multi-media devices to move and change objects in space
 - transformation geometry
- * We only consider solving equations on Maths1 with using 2×2 matrices

- * We can use our 2x2 matrices to express 2 simultaneous equations (2 equations about the same 2 variables)
- * First we must put them in the correct format
- * for the variables x & y the format should be

$$\begin{aligned} ax + by &= m \\ cx + dy &= n \end{aligned} \quad \{\text{where } a, b, c, d, m \text{ \& } n \text{ are constants}\}$$

Example

Peter and Jane spend £240 altogether and Peter spends 3 times as much as Jane.

let p : what Peter spends and j : what Jane spends

then $p + j = 240$	(right format a and $b = 1$ $m = 240$)
$p = 3j$	(wrong format)
rewrite $p - 3j = 0$	(right format $c = 1$ $d = -3$ $n = 0$)

We can use our 2x2 matrices to express these simultaneous equations

$$\left. \begin{array}{l} x + y = 240 \\ x - 3y = 0 \end{array} \right\}$$

Becomes in matrix form

$$\begin{pmatrix} 1 & 1 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 240 \\ 0 \end{pmatrix}$$

constants
from the
left hand
side

UNKNOWNNS
X ~ x1
Y ~ x2

constants
from the
right hand
side

$$\begin{pmatrix} 1 & 1 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 240 \\ 0 \end{pmatrix}$$

Format is $Ax=B$

To solve this using the matrix we must get rid of it by using its inverse!

First find the inverse

$$\begin{pmatrix} 1 & 1 \\ 1 & -3 \end{pmatrix}^{-1} = \frac{1}{-4} \begin{pmatrix} -3 & -1 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} 0.75 & 0.25 \\ 0.25 & -0.25 \end{pmatrix}$$

now use it on both sides of the equation

$$\begin{pmatrix} 0.75 & 0.25 \\ 0.25 & -0.25 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & -3 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0.75 & 0.25 \\ 0.25 & -0.25 \end{pmatrix} \begin{pmatrix} 240 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 0.75 & 0.25 \\ 0.25 & -0.25 \end{pmatrix} \begin{pmatrix} 240 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = \begin{pmatrix} 180 \\ 60 \end{pmatrix}$$

So Answer is $p = 180$ $j = 60$

1. Format the simultaneous equations for variable x & y

$$ax + by = m$$

$$cx + dy = n$$

2. Rewrite them in matrix form

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} m \\ n \end{pmatrix}$$

3. Find the inverse of the 2×2 matrix

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$

4. Solve for the variables x, y by multiplying the right hand side of the equation by the inverse

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} \begin{pmatrix} m \\ n \end{pmatrix}$$

$$3x + 4y = 5$$

$$5x = 7 - 6y$$

$$\begin{pmatrix} 3 & 4 \\ 5 & 6 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 5 \\ 7 \end{pmatrix}$$

$$\text{Answer } x = -1 \quad y = 2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{-2} \begin{pmatrix} 6 & -4 \\ -5 & 3 \end{pmatrix} \begin{pmatrix} 5 \\ 7 \end{pmatrix} = \frac{1}{-2} \begin{pmatrix} 30 - 28 \\ -25 + 21 \end{pmatrix} = \frac{1}{-2} \begin{pmatrix} 2 \\ -4 \end{pmatrix} = \begin{pmatrix} -1 \\ 2 \end{pmatrix}$$

$$x + 7y = 1.24$$

$$3y - x = 0.76$$

$$\begin{pmatrix} 1 & 7 \\ -1 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1.24 \\ 0.76 \end{pmatrix}$$

$$\text{Answer } x = -0.16 \quad y = 0.2$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{10} \begin{pmatrix} 3 & -7 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1.24 \\ 0.76 \end{pmatrix} = \frac{1}{10} \begin{pmatrix} 3.72 - 5.32 \\ 1.24 + 0.76 \end{pmatrix} = \frac{1}{10} \begin{pmatrix} -1.6 \\ 2 \end{pmatrix} = \begin{pmatrix} -0.16 \\ 0.2 \end{pmatrix}$$

$$8x = 3y - 1$$

$$x + y = -7$$

$$\begin{pmatrix} 8 & -3 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} -1 \\ -7 \end{pmatrix}$$

$$\text{Answer } x = -2 \quad y = -5$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{11} \begin{pmatrix} 1 & 3 \\ -1 & 8 \end{pmatrix} \begin{pmatrix} -1 \\ -7 \end{pmatrix} = \frac{1}{11} \begin{pmatrix} -1 - 21 \\ 1 - 56 \end{pmatrix} = \frac{1}{11} \begin{pmatrix} -22 \\ -55 \end{pmatrix} = \begin{pmatrix} -2 \\ -5 \end{pmatrix}$$