

Volume Integral


Scalar Field, F Integral

If V is a closed region and F is a scalar field in region V , volume integral F of V is

$$\int_V F dV = \iiint_V F dx dy dz$$

Example :

Scalar function $F = 2x$ defined in one cubic that has been built by planes $x = 0$, $x = 1$, $y = 0$, $y = 3$, $z = 0$ and $z = 2$. Evaluate volume integral F of the cubic.



Solution

$$\begin{aligned}\int_V F dV &= \int_{z=0}^2 \int_{y=0}^3 \int_{x=0}^1 2x dx dy dz \\ &= 2 \int_{z=0}^2 \int_{y=0}^3 \left[\frac{x^2}{2} \right]_0^1 dy dz \\ &= 2 \int_{z=0}^2 \int_{y=0}^3 \frac{1}{2} dy dz \\ &= 2 \cdot \frac{1}{2} \int_{z=0}^2 [y]_0^3 dz \\ &= \int_{z=0}^2 3 dz = 3[z]_0^2 = 6\end{aligned}$$

Volume Integral

If V is a closed region and \vec{F} , vector field in region V , Volume integral \vec{F} of V is


$$\int_V \vec{F} dV = \int_{x_1}^{x_2} \int_{y_1}^{y_2} \int_{z_1}^{z_2} \vec{F} dz dy dx$$

Example:

Evaluate $\int_V \vec{F} dV$, where V is a region bounded by $x = 0$, $y = 0$, $z = 0$ and $2x + y + z = 2$, and also given $\vec{F} = 2z \vec{i} + y \vec{k}$

Solution

If $x = y = 0$, plane $2x + y + z = 2$ intersects z -axis at $z = 2$.


 $(0,0,2)$

If $x = z = 0$, plane $2x + y + z = 2$ intersects y -axis at $y = 2$.

 $(0,2,0)$

If $y = z = 0$, plane $2x + y + z = 2$ intersects x -axis at $x = 1$.

 $(1,0,0)$



We can generate this integral in 3 steps :

1. Line Integral from $x = 0$ to $x = 1$.
2. Surface Integral from line $y = 0$ to line $y = 2(1-x)$.
3. Volume Integral from surface $z = 0$ to surface $2x + y + z = 2$ that is $z = 2(1-x) - y$

Therefore,


$$\begin{aligned}\int_V \vec{F} dV &= \int_{x=0}^1 \int_{y=0}^{2(1-x)} \int_{z=0}^{2(1-x)-y} \vec{F} dz dy dx \\ &= \int_{x=0}^1 \int_{y=0}^{2(1-x)} \int_{z=0}^{2(1-x)-y} (2z \vec{i} + y \vec{k}) dz dy dx \\ &\quad \vdots \\ &\quad \vdots \\ &= \frac{2}{3} \vec{i} + \frac{1}{3} \vec{k}\end{aligned}$$

Example:

Evaluate $\int_V \vec{F} dV$ where $\vec{F} = 2\vec{i} + 2z\vec{j} + y\vec{k}$

and V is region bounded by $z = 0$, $z = 4$ and

$$x^2 + y^2 = 9$$



Using polar coordinate of cylinder,

$$x = \rho \cos \phi; \quad y = \rho \sin \phi; \quad z = z;$$

$$dV = \rho d\rho d\phi dz$$

where

$$0 \leq \rho \leq 3, \quad 0 \leq \phi \leq 2\pi, \quad 0 \leq z \leq 4$$

Therefore,

$$\begin{aligned}\int_V \underline{F} dV &= \iiint_V (2\underline{i} + 2z\underline{j} + y\underline{k}) dx dy dz \\ &= \int_{z=0}^4 \int_{\phi=0}^{2\pi} \int_{\rho=0}^3 (2\underline{i} + 2z\underline{j} + \rho \sin \phi \underline{k}) \rho d\rho d\phi dz \\ &\quad \vdots \\ &\quad \vdots \\ &= 72\pi \underline{i} + 144\pi \underline{j}\end{aligned}$$

Exercise:

Evaluate $\int_V \vec{F} dV$ where $\vec{F} = 3\vec{i} + z\vec{j} + 2y\vec{k}$ and

V is region bounded by planes $z = 0$, $z = 3$
and surface $x^2 + y^2 = 4$.

Answer: $18\pi \left(2\vec{i} + \vec{j} \right)$