

Introduction to Electromagnetic Fields;
Maxwell's Equations;
Electromagnetic Fields in Materials;




- To provide an overview of classical electromagnetics, Maxwell's equations, electromagnetic fields in materials, and phasor concepts.
- To begin our study of electrostatics with Coulomb's law; definition of electric field; computation of electric field from discrete and continuous charge distributions; and scalar electric potential.

Introduction to Electromagnetic Fields

- **Electromagnetics** is the study of the effect of charges at rest and charges in motion.
- Some special cases of electromagnetics:
 - **Electrostatics**: charges at rest
 - **Magnetostatics**: charges in steady motion (DC)
 - **Electromagnetic waves**: waves excited by charges in time-varying motion

Introduction to Electromagnetic Fields




Introduction to Electromagnetic Fields



- transmitter and receiver are connected by a “field.”


Introduction to Electromagnetic Fields

High-speed, high-density digital circuits:



- consider an interconnect between points “1” and “2”

Introduction to Electromagnetic Fields



- Propagation delay
- Electromagnetic coupling
- Substrate modes

Introduction to Electromagnetic Fields

- When an event in one place has an effect on something at a different location, we talk about the events as being connected by a “field”.
- A *field* is a spatial distribution of a quantity; in general, it can be either *scalar* or *vector* in nature.

Introduction to Electromagnetic Fields

- Electric and magnetic fields:
 - Are vector fields with three spatial components.
 - Vary as a function of position in 3D space as well as time.
 - Are governed by partial differential equations derived from Maxwell's equations.

Introduction to Electromagnetic Fields

- A *scalar* is a quantity having only an amplitude (and possibly phase).

Examples: voltage, current, charge, energy, temperature

- A *vector* is a quantity having direction in addition to amplitude (and possibly phase).

Examples: velocity, acceleration, force

Introduction to Electromagnetic Fields

➤ Fundamental vector field quantities in electromagnetics:

- Electric field intensity (\underline{E})
units = volts per meter ($V/m = kg\ m/A/s^3$)
- Electric flux density (electric displacement) (\underline{D})
units = coulombs per square meter ($C/m^2 = A\ s/m^2$)
- Magnetic field intensity (\underline{H})
units = amps per meter (A/m)
- Magnetic flux density (\underline{B})
units = teslas = webers per square meter ($T = Wb/m^2 = kg/A/s^3$)

Introduction to Electromagnetic Fields

- Universal constants in electromagnetics:
 - Velocity of an electromagnetic wave (e.g., light) in free space (perfect vacuum)

$$c \approx 3 \times 10^8 \text{ m/s}$$

- Permeability of free space

$$\mu_0 = 4\pi \times 10^{-7} \text{ H/m}$$

- Permittivity of free space:

$$\epsilon_0 \approx 8.854 \times 10^{-12} \text{ F/m}$$

- Intrinsic impedance of free space:

$$\eta_0 \approx 120\pi \Omega$$

Introduction to Electromagnetic Fields

- Relationships involving the universal constants:


$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}} \quad \eta_0 = \sqrt{\frac{\mu_0}{\epsilon_0}}$$

In free space:

$$\underline{B} = \mu_0 \underline{H}$$

$$\underline{D} = \epsilon_0 \underline{E}$$

Introduction to Electromagnetic Fields



Maxwell's Equations

- *Maxwell's equations in integral form* are the fundamental postulates of classical electromagnetics - all classical electromagnetic phenomena are explained by these equations.
- Electromagnetic phenomena include electrostatics, magnetostatics, electromagnetostatics and electromagnetic wave propagation.
- The differential equations and boundary conditions that we use to formulate and solve EM problems are all derived from *Maxwell's equations in integral form*.

Maxwell's Equations

- Various *equivalence principles* consistent with Maxwell's equations allow us to replace more complicated electric current and charge distributions with *equivalent magnetic sources*.
- These *equivalent magnetic sources* can be treated by a generalization of Maxwell's equations.

Maxwell's Equations in Integral Form (Generalized to Include Equivalent Magnetic Sources)

$$\oint_C \underline{E} \cdot d\underline{l} = -\frac{d}{dt} \int_S \underline{B} \cdot d\underline{S} - \int_S \underline{K}_c \cdot d\underline{S} - \int_S \underline{K}_i \cdot d\underline{S}$$

$$\oint_C \underline{H} \cdot d\underline{l} = \frac{d}{dt} \int_S \underline{D} \cdot d\underline{S} + \int_S \underline{J}_c \cdot d\underline{S} + \int_S \underline{J}_i \cdot d\underline{S}$$

$$\oint_S \underline{D} \cdot d\underline{S} = \int_V q_{ev} dv$$

$$\oint_S \underline{B} \cdot d\underline{S} = \int_V q_{mv} dv$$

Adding the fictitious magnetic source terms is equivalent to living in a universe where magnetic monopoles (charges) exist.


Continuity Equation in Integral Form (Generalized to Include Equivalent Magnetic Sources)

$$\oint_S \underline{J} \cdot d\underline{s} = -\frac{\partial}{\partial t} \int_V q_{ev} dv$$


$$\oint_S \underline{K} \cdot d\underline{s} = -\frac{\partial}{\partial t} \int_V q_{mv} dv$$

- The *continuity equations* are implicit in Maxwell's equations.

Contour, Surface and Volume Conventions



- open surface S bounded by closed contour C
- $d\underline{S}$ in direction given by RH rule



- volume V bounded by closed surface S
- $d\underline{S}$ in direction outward from V

Electric Current and Charge Densities

- $J_c =$ (electric) conduction current density (A/m^2)
- $J_i =$ (electric) impressed current density (A/m^2)
- $q_{ev} =$ (electric) charge density (C/m^3)

Magnetic Current and Charge Densities

- K_c = magnetic conduction current density (V/m^2)
- K_i = magnetic impressed current density (V/m^2)
- q_{mv} = magnetic charge density (Wb/m^3)

Maxwell's Equations in Differential Form (Generalized to Include Equivalent Magnetic Sources)

$$\nabla \times \underline{E} = -\frac{\partial \underline{B}}{\partial t} - \underline{K}_c - \underline{K}_i$$

$$\nabla \times \underline{H} = \frac{\partial \underline{D}}{\partial t} + \underline{J}_c + \underline{J}_i$$

$$\nabla \cdot \underline{D} = q_{ev}$$

$$\nabla \cdot \underline{B} = q_{mv}$$

Continuity Equation in Differential Form (Generalized to Include Equivalent Magnetic Sources)

$$\nabla \cdot \underline{J} = -\frac{\partial q_{ev}}{\partial t}$$

$$\nabla \cdot \underline{K} = -\frac{\partial q_{mv}}{\partial t}$$

- The *continuity equations* are implicit in Maxwell's equations.

Electromagnetic Fields in Materials

➤ In free space, we have:

$$\underline{D} = \varepsilon_0 \underline{E}$$

$$\underline{B} = \mu_0 \underline{H}$$

$$\underline{J}_c = 0$$

$$\underline{K}_c = 0$$

Electromagnetic Fields in Materials

➤ In a *simple medium*, we have:

$$\underline{D} = \varepsilon \underline{E}$$

$$\underline{B} = \mu \underline{H}$$

$$\underline{J}_c = \sigma \underline{E}$$

$$\underline{K}_c = \sigma_m \underline{H}$$

- *linear* (independent of field strength)
- *isotropic* (independent of position within the medium)
- *homogeneous* (independent of direction)
- *time-invariant* (independent of time)
- *non-dispersive* (independent of frequency)

Electromagnetic Fields in Materials

- $\epsilon = \text{permittivity} = \epsilon_r \epsilon_0$ (F/m)
- $\mu = \text{permeability} = \mu_r \mu_0$ (H/m)
- $\sigma = \text{electric conductivity} = \epsilon_r \epsilon_0$ (S/m)
- $\sigma_m = \text{magnetic conductivity} = \epsilon_r \epsilon_0$ (Ω/m)

Maxwell's Equations in Differential Form for Time-Harmonic Fields in Simple Medium


$$\nabla \times \underline{E} = -(j\omega\mu + \sigma_m) \underline{H} - \underline{K}_i$$

$$\nabla \times \underline{H} = (j\omega\varepsilon + \sigma) \underline{E} + \underline{J}_i$$

$$\nabla \cdot \underline{E} = \frac{q_{ev}}{\varepsilon}$$

$$\nabla \cdot \underline{H} = \frac{q_{mv}}{\mu}$$

Electrostatics as a Special Case of Electromagnetics



Electrostatics

- *Electrostatics* is the branch of electromagnetics dealing with the effects of electric charges at rest.
- The fundamental law of *electrostatics* is *Coulomb's law*.

Electric Charge

- Electrical phenomena caused by friction are part of our everyday lives, and can be understood in terms of *electrical charge*.
- The effects of *electrical charge* can be observed in the attraction/repulsion of various objects when “charged.”
- Charge comes in two varieties called “positive” and “negative.”

Electric Charge

- Objects carrying a net positive charge attract those carrying a net negative charge and repel those carrying a net positive charge.
- Objects carrying a net negative charge attract those carrying a net positive charge and repel those carrying a net negative charge.
- On an atomic scale, electrons are negatively charged and nuclei are positively charged.

Continuous Distributions of Charge


➤ Charge can occur as

- *point charges* (C)
- *volume charges* (C/m³)
- *surface charges* (C/m²)
- *line charges* (C/m)

⇐ **most general**

Continuous Distributions of Charge


➤ Volume charge density



$$q_{ev}(\underline{r}') = \lim_{\Delta V' \rightarrow 0} \frac{Q_{encl}}{\Delta V'}$$

Continuous Distributions of Charge

- Electric field due to volume charge density




$$d\underline{E}(\underline{r}) = \frac{q_{ev}(\underline{r}')dV' \underline{R}}{4\pi\epsilon_0 R^3}$$

Electric Field Due to Volume Charge Density

$$\underline{E}(\underline{r}) = \frac{1}{4\pi\epsilon_0} \int_{V'} \frac{q_{ev}(\underline{r}') \underline{R}}{R^3} dv'$$

Continuous Distributions of Charge


➤ Surface charge density



$$q_{es}(\underline{r}') = \lim_{\Delta S' \rightarrow 0} \frac{Q_{encl}}{\Delta S'}$$

Continuous Distributions of Charge

- Electric field due to surface charge density




$$d\underline{E}(\underline{r}) = \frac{q_{es}(\underline{r}')ds' \underline{R}}{4\pi\epsilon_0 R^3}$$

Electric Field Due to Surface Charge Density

$$\underline{E}(\underline{r}) = \frac{1}{4\pi\epsilon_0} \int_{S'} \frac{q_{es}(\underline{r}') \underline{R}}{R^3} ds'$$

Continuous Distributions of Charge


➤ Line charge density



$$q_{el}(\underline{r}') = \lim_{\Delta L' \rightarrow 0} \frac{Q_{encl}}{\Delta L'}$$

Continuous Distributions of Charge

- Electric field due to line charge density



$$d\underline{E}(\underline{r}) = \frac{q_{el}(\underline{r}')d\underline{l}' \underline{R}}{4\pi\epsilon_0 R^3}$$

Electric Field Due to Line Charge Density

$$\underline{E}(\underline{r}) = \frac{1}{4\pi\epsilon_0} \int_{L'} \frac{q_{el}(\underline{r}') \underline{R}}{R^3} dl'$$